

Constructing new optimal entanglement witnesses

Dariusz Chruściński, Justyna Pytel and Gniewomir Sarbicki
*Institute of Physics, Nicolaus Copernicus University,
Grudziądzka 5/7, 87-100 Toruń, Poland*

We provide a new class of indecomposable entanglement witnesses. In 4×4 case it reproduces the well known Breuer-Hall witness. We prove that these new witnesses are optimal and atomic, i.e. they are able to detect the “weakest” quantum entanglement encoded into states with positive partial transposition (PPT). Equivalently, we provide a new construction of indecomposable atomic maps in the algebra of $2k \times 2k$ complex matrices. It is shown that their structural physical approximations give rise to entanglement breaking channels. This result supports recent conjecture by Korbicz *et al.*

PACS numbers: 03.65.Ud, 03.67.-a

I. INTRODUCTION

The interest on quantum entanglement has dramatically increased during the last two decades due to the emerging field of quantum information theory [1]. It turns out that quantum entanglement may be used as basic resources in quantum information processing and communication. The prominent examples are quantum cryptography, quantum teleportation, quantum error correction codes and quantum computation.

Since the quantum entanglement is the basic resource for the new quantum information technologies it is therefore clear that there is a considerable interest in efficient theoretical and experimental methods of entanglement detection (see [2] for the review).

The most general approach to characterize quantum entanglement uses a notion of an entanglement witness (EW) [3, 4]. A Hermitian operator W defined on a tensor product $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$ is called an EW iff 1) $\text{Tr}(W\sigma_{\text{sep}}) \geq 0$ for all separable states σ_{sep} , and 2) there exists an entangled state ρ such that $\text{Tr}(W\rho) < 0$ (one says that ρ is detected by W). It turns out that a state is entangled if and only if it is detected by some EW [3]. There was a considerable effort in constructing and analyzing the structure of EWs [5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15]. In fact, entanglement witnesses have been measured in several experiments [16, 17]. Moreover, several procedures for optimizing EWs for arbitrary states were proposed [6, 18, 19, 20]. It should be stressed that there is no universal W , i.e. there is no entanglement witness which detects all entangled states. Each entangled state ρ may be detected by a specific choice of W . It is clear that each EW provides a new separability test and it may be interpreted as a new type of Bell inequality [21]. There is, however, no general procedure for constructing EWs.

Due to the Choi-Jamiołkowski isomorphism [22, 23] any EW corresponds to a linear positive map $\Lambda : \mathcal{B}(\mathcal{H}_A) \rightarrow \mathcal{B}(\mathcal{H}_B)$, where by $\mathcal{B}(\mathcal{H})$ we denote the space

of bounded operators on the Hilbert space $\mathcal{H}$. Recall that a linear map Λ is said to be positive if it sends a positive operator on $\mathcal{H}_A$ into a positive operator on $\mathcal{H}_B$. It turns out [3] that a state ρ in $\mathcal{H}_A \otimes \mathcal{H}_B$ is separable iff $(\mathbb{1}_A \otimes \Lambda)\rho$ is positive definite for all positive maps $\Lambda : \mathcal{B}(\mathcal{H}_B) \rightarrow \mathcal{B}(\mathcal{H}_A)$ (actually this result is based on [24]). Unfortunately, in spite of the considerable effort, the structure of positive maps is rather poorly understood [24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41].

In the present paper we provide a construction of a new class of positive maps in $\mathcal{B}(\mathbb{C}^{2k})$ with $k \geq 2$. Our construction uses the well-known reduction map as a building block. It turns out that for $k = 2$ our construction reproduces Breuer-Hall maps [36, 37] but for $k > 2$ it gives completely new family of maps. It is shown that proposed maps are indecomposable (i.e. they are able to detect entangled PPT states) and atomic (i.e. they are able to detect “weakly” entangled PPT states). As a byproduct we construct new families of PPT entangled states detected by our maps.

The paper is organized as follows: for pedagogical reason we collect basic definitions and introduce the most important properties of positive maps and entanglement witnesses in Section II. Section III provides basic construction. Then in Section IV we study basic properties of our maps/witnesses (indecomposability, atomicity, optimality). Section V discusses structural physical approximation (SPA) [42, 43, 44] of our maps. It is shown that the corresponding SPA gives rise to entanglement breaking channels and hence it supports recent conjecture by Korbicz *et al.* [44]. Final conclusions are collected in the last Section.

II. POSITIVE MAPS, ENTANGLEMENT WITNESSES AND ALL THAT

For the reader convenience we recall basic definitions and properties which are important throughout this paper.

A. Positive maps

Let $\Lambda : \mathcal{B}(\mathcal{H}_A) \rightarrow \mathcal{B}(\mathcal{H}_B)$ be a positive linear map. In what follows we shall consider only finite dimensional Hilbert spaces such that $\dim \mathcal{H}_A = d_A$ and $\dim \mathcal{H}_B = d_B$. One calls Λ k -positive if

$$\mathbb{1}_k \otimes \Lambda : M_k \otimes \mathcal{B}(\mathcal{H}_A) \longrightarrow M_k \otimes \mathcal{B}(\mathcal{H}_B) , \quad (1)$$

is positive. In the above formula M_k denotes a linear space of $k \times k$ complex matrices and $\mathbb{1}_k : M_k \rightarrow M_k$ is an identity map, i.e. $\mathbb{1}_k(A) = A$ for each $A \in M_k$. A positive map which is k -positive for each k is called completely positive (CP). Actually, if $d_A, d_B < \infty$ one shows [22] that Λ is CP iff it is d -positive with $d = \min\{d_A, d_B\}$.

Definition 1 A positive map Λ is decomposable if

$$\Lambda = \Lambda_1 + \Lambda_2 \circ T , \quad (2)$$

where Λ_1 and Λ_2 are CP and T denotes transposition in a given basis. Maps which are not decomposable are called indecomposable (or nondecomposable).

Definition 2 A positive map Λ is atomic if it cannot be represented as

$$\Lambda = \Lambda_1 + \Lambda_2 \circ T , \quad (3)$$

where Λ_1 and Λ_2 are 2-positive.

Definition 3 A positive map Λ is optimal if and only if for any CP map Φ , the map $\Lambda - \Phi$ is no longer positive.

B. Entanglement witnesses

Using Choi-Jamiołkowski isomorphism [22, 23] each positive map Λ gives rise to entanglement witness W

$$W = d_A(\mathbb{1}_A \otimes \Lambda)P_A^+ , \quad (4)$$

where P_A^+ denotes maximally entangled state in $\mathbb{C}^{d_A} \otimes \mathbb{C}^{d_A}$ and $\mathbb{1}_A$ denotes an identity map acting on $\mathcal{B}(\mathcal{H}_A)$. One has an obvious

Definition 4 An entanglement witness W defined by (4) is decomposable/indecomposable (atomic) [optimal] $\{k$ -EW $\}$ if and only if the corresponding positive map Λ is decomposable/indecomposable (atomic) [optimal] $\{k$ -positive $\}$.

It is clear that $W \in \mathcal{B}(\mathcal{H}_A \otimes \mathcal{H}_B)$ is a decomposable EW iff

$$W = A + B^\Gamma , \quad (5)$$

where $A, B \geq 0$ and $B^\Gamma = (\mathbb{1}_A \otimes T)B$ denotes partial transposition. Witnesses which cannot be represented as in (5) are indecomposable

Let ψ be a normalize vector in $\mathcal{H}_A \otimes \mathcal{H}_B$. Denote by $\text{SR}(\psi)$ the number of nonvanishing Schmidt coefficients of ψ . One has

$$1 \leq \text{SR}(\psi) \leq d . \quad (6)$$

Now, W is k -EW iff

$$\langle \psi | W | \psi \rangle \geq 0 , \quad (7)$$

for each ψ such that $\text{SR}(\psi) \leq k$. Evidently, $W \geq 0$ iff W is d -EW. Now, W is atomic if it cannot be represented as

$$W = W_1 + W_2^\Gamma , \quad (8)$$

where W_1 and W_2 are 2-EWs. Finally, W is optimal EW iff for any $P \geq 0$, $W - P$ is no longer EW. Following [6] one has the following criterion for the optimality of W : if the set of product vectors $\psi \otimes \phi \in \mathcal{H}_A \otimes \mathcal{H}_B$ satisfying

$$\langle \psi \otimes \phi | W | \psi \otimes \phi \rangle = 0 , \quad (9)$$

span the total Hilbert space $\mathcal{H}_A \otimes \mathcal{H}_B$, then W is optimal.

C. Detecting quantum entanglement

Positive maps and EWs are basic tools in detecting quantum entanglement. A state ρ in $\mathcal{H}_A \otimes \mathcal{H}_B$ is separable if and only if for all positive maps $\Lambda : \mathcal{B}(\mathcal{H}_B) \rightarrow \mathcal{B}(\mathcal{H}_A)$ one has

$$(\mathbb{1}_A \otimes \Lambda)\rho \geq 0 . \quad (10)$$

Equivalently, iff for each entanglement witness W

$$\text{Tr}(\rho W) \geq 0 . \quad (11)$$

Note that entangled PPT states can be detected by indecomposable maps/witnesses only. Let σ be a density operator in $\mathcal{H}_A \otimes \mathcal{H}_B$. Following [45] one introduces its Schmidt number

$$\text{SN}(\sigma) = \min_{p_k, \psi_k} \left\{ \max_k \text{SR}(\psi_k) \right\} , \quad (12)$$

where the minimum is taken over all possible pure states decompositions

$$\sigma = \sum_k p_k |\psi_k\rangle\langle\psi_k|, \quad (13)$$

with $p_k \geq 0$, $\sum_k p_k = 1$ and ψ_k are normalized vectors in $\mathcal{H}_A \otimes \mathcal{H}_B$. Note, that if $\sigma = |\psi\rangle\langle\psi|$, then $\text{SN}(\sigma) = \text{SR}(\psi)$. Again one has $1 \leq \text{SN}(\sigma) \leq d$. Suppose now that σ is PPT but entangled. Intuitively, the ‘weakest’ quantum entangled encoded in σ corresponds to the situation when $\text{SR}(\sigma) = \text{SR}(\sigma^\Gamma) = 2$. Such ‘weakly’ entangled PPT states can be detected by atomic maps/witnesses only.

III. REDUCTION MAP AS A BUILDING BLOCK

Let us start with an elementary positive map in $\mathcal{B}(\mathbb{C}^n)$ called reduction map [46]

$$R_n(X) = \mathbb{I}_n \text{Tr} X - X, \quad (14)$$

for $X \in \mathcal{B}(\mathbb{C}^n)$. It is well known that R_n is completely co-positive (i.e. $R_n \circ T$ is CP) and hence optimal. Recently, this map was generalized by Breuer and Hall [36, 37] to the following family of positive maps

$$\Phi_{2k}^U(X) = \frac{1}{2(k-1)} (R_{2k}(X) - UX^T U^\dagger), \quad (15)$$

where U is an arbitrary antisymmetric unitary $2k \times 2k$ matrix. It was shown that these maps are indecomposable [36, 37] and optimal [36]. Such antisymmetric unitary matrix may be easily constructed as follows

$$U = V U_0 V^\dagger, \quad (16)$$

where V stands for real orthogonal matrix ($VV^\dagger = VV^T = \mathbb{I}_{2k}$) and

$$U_0 = \mathbb{I}_k \otimes J, \quad (17)$$

with J being 2×2 symplectic matrix

$$J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (18)$$

It is therefore clear that in this case one has

$$\Phi_{2k}^U(X) = V \Phi_{2k}^0 (V^\dagger X V) V^\dagger, \quad (19)$$

where Φ_{2k}^0 corresponds to Φ_{2k}^U with $U = U_0$. Actually, one can always find a basis in $\mathbb{C}^{2k}$ such that U takes the ‘canonical form’ U_0 . Interestingly for $k = 2$ the Breuer-Hall map Φ_4^0 reproduces well known Robertson map [28] who provided it as an example of an extremal (and hence optimal) indecomposable positive map. Moreover, Robertson construction may be nicely described in terms of R_2 as follows [39]

$$\Phi_4^0 \left(\begin{array}{c|c} X_{11} & X_{12} \\ \hline X_{21} & X_{22} \end{array} \right) = \frac{1}{2} \left(\begin{array}{c|c} \mathbb{I}_2 \text{Tr} X_{22} & -[X_{12} + R_2(X_{21})] \\ \hline -[X_{21} + R_2(X_{12})] & \mathbb{I}_2 \text{Tr} X_{11} \end{array} \right), \quad (20)$$

where $X_{kl} \in \mathcal{B}(\mathbb{C}^2)$. This pattern is reproduced for arbitrary k . It is easy to show that the action of Φ_{2k}^0

may be represented as follows:

$$\Phi_{2k}^0 \left(\begin{array}{c|c|c|c} X_{11} & X_{12} & \cdots & X_{1k} \\ \hline X_{21} & X_{22} & \cdots & X_{2k} \\ \hline \vdots & \vdots & \ddots & \vdots \\ \hline X_{k1} & X_{k2} & \cdots & X_{kk} \end{array} \right) = \frac{1}{2(k-1)} \left(\begin{array}{c|c|c|c} \mathbb{I}_2(\text{Tr} X - \text{Tr} X_{11}) & -(X_{12} + R_2(X_{21})) & \cdots & -(X_{1k} + R_2(X_{k1})) \\ \hline -(X_{21} + R_2(X_{12})) & \mathbb{I}_2(\text{Tr} X - \text{Tr} X_{22}) & \cdots & -(X_{2k} + R_2(X_{k2})) \\ \hline \vdots & \vdots & \ddots & \vdots \\ \hline -(X_{k1} + R_2(X_{1k})) & -(X_{k2} + R_2(X_{2k})) & \cdots & \mathbb{I}_2(\text{Tr} X - \text{Tr} X_{kk}) \end{array} \right),$$

where again X_{ij} are 2×2 blocks. Hence Φ_{2k}^0 is defined in (15) by R_{2k} but the above pattern shows that it

basically uses reduction map R_2 only. We stress that R_2 is exceptional: it is not only optimal but also ex-

tremal. Indeed, the corresponding entanglement witness $W_2 = 2(\mathbb{I} \otimes R_2)P_2^+$ reads as follows

$$W_2 = \mathbb{I}_2 \otimes \mathbb{I}_2 - P_2^+ = \left(\begin{array}{cc|cc} \cdot & \cdot & \cdot & -1 \\ \cdot & 1 & \cdot & \cdot \\ \hline \cdot & \cdot & 1 & \cdot \\ -1 & \cdot & \cdot & \cdot \end{array} \right). \quad (21)$$

and $W_2 = P^\Gamma$, where $P = |\psi\rangle\langle\psi|$ with

$$|\psi\rangle = |01\rangle - |10\rangle. \quad (22)$$

Note, that R_2 may be nicely represented as follows

$$R_2(X) = JX^T J^\dagger, \quad (23)$$

with J defined in (18). Formula (23) provides Kraus representation for $R_2 \circ T$ and shows that R_2 is completely co-positive.

In the present paper we propose another construction of maps in $\mathcal{B}(\mathbb{C}^{2k})$. Now, instead of treating a $2k \times 2k$ matrix X as a $k \times k$ matrix with 2×2 blocks X_{ij} we consider alternative possibility, i.e. we consider X as a 2×2 with $k \times k$ blocks and define

$$\Psi_{2k}^0 \left(\begin{array}{cc|cc} X_{11} & X_{12} & & \\ X_{21} & X_{22} & & \end{array} \right) = \frac{1}{k} \left(\begin{array}{cc|cc} \mathbb{I}_k \text{Tr} X_{22} & & -[X_{12} + R_k(X_{21})] & \\ & -[X_{21} + R_k(X_{12})] & & \mathbb{I}_k \text{Tr} X_{11} \end{array} \right). \quad (24)$$

Again, normalization factor guaranties that the map is unital, i.e. $\Psi_{2k}^0(\mathbb{I}_2 \otimes \mathbb{I}_k) = \mathbb{I}_2 \otimes \mathbb{I}_k$. It is clear that for $k = 2$ one has

$$\Phi_4^0 = \Psi_4^0. \quad (25)$$

We stress that our new construction is much simpler than Φ_{2k}^0 and it uses as a building block the true reduction map in $\mathcal{B}(\mathbb{C}^k)$. Moreover, it is clear that it provides a natural generalization of the original Robertson map in $\mathcal{B}(\mathbb{C}^4)$.

Now, our task is to prove that Ψ_{2k}^0 defines a positive map. It is enough to show that each rank-1 projector P is mapped via Ψ_{2k}^0 into a positive element in $\mathcal{B}(\mathbb{C}^{2k})$, that is, $\Psi_{2k}^0(P) \geq 0$. Let $P = |\psi\rangle\langle\psi|$ with arbitrary ψ from $\mathbb{C}^{2k}$. Now, due to $\mathbb{C}^{2k} = \mathbb{C}^k \oplus \mathbb{C}^k$ one has

$$\psi = \psi_1 \oplus \psi_2, \quad (26)$$

with $\psi_1, \psi_2 \in \mathbb{C}^k$ and hence

$$P = \left(\begin{array}{cc|cc} X_{11} & X_{12} & & \\ X_{21} & X_{22} & & \end{array} \right) = \left(\begin{array}{cc|cc} |\psi_1\rangle\langle\psi_1| & |\psi_1\rangle\langle\psi_2| & & \\ |\psi_2\rangle\langle\psi_1| & |\psi_2\rangle\langle\psi_2| & & \end{array} \right). \quad (27)$$

One has therefore

$$\Psi_{2k}^0(P) = \frac{1}{k} \left(\begin{array}{cc|cc} \mathbb{I}_k \text{Tr} X_{22} & & -A & \\ & -A^\dagger & & \mathbb{I}_k \text{Tr} X_{11} \end{array} \right), \quad (28)$$

where the linear operator $A : \mathbb{C}^k \rightarrow \mathbb{C}^k$ reads as follows

$$A = |\psi_1\rangle\langle\psi_2| - |\psi_2\rangle\langle\psi_1| + \langle\psi_1|\psi_2\rangle \mathbb{I}_k. \quad (29)$$

Let $\langle\psi_j|\psi_j\rangle = a_j^2 > 0$ (if one of a_j vanishes then evidently one has $\Psi_{2k}^0(P) \geq 0$). Defining

$$L = \sqrt{k} \left(\begin{array}{cc|cc} \mathbb{I}_k a_2^{-1} & & \mathbb{O}_k & \\ & \mathbb{O}_k & & \mathbb{I}_k a_1^{-1} \end{array} \right), \quad (30)$$

one finds

$$L \Psi_{2k}^0(P) L^\dagger = \left(\begin{array}{cc|cc} \mathbb{I}_k & & -\tilde{A} & \\ & -\tilde{A}^\dagger & & \mathbb{I}_k \end{array} \right), \quad (31)$$

with

$$\tilde{A} = |\tilde{\psi}_1\rangle\langle\tilde{\psi}_2| - |\tilde{\psi}_2\rangle\langle\tilde{\psi}_1| + \langle\tilde{\psi}_1|\tilde{\psi}_2\rangle \mathbb{I}_k, \quad (32)$$

and normalized $\tilde{\psi}_j = \psi_j/a_j$. Hence, to show that $\Psi_{2k}^0(P) \geq 0$ one needs to prove

$$\left(\begin{array}{cc|cc} \mathbb{I}_k & & -\tilde{A} & \\ & -\tilde{A}^\dagger & & \mathbb{I}_k \end{array} \right) \geq 0, \quad (33)$$

for arbitrary $\psi_j \neq 0$. Now, the above condition is equivalent to

$$\tilde{A} \tilde{A}^\dagger \leq \mathbb{I}_k. \quad (34)$$

Vectors $\{\psi_1, \psi_2\}$ span 2-dimensional subspace in $\mathbb{C}^k$ and let $\{e_1, e_2\}$ be a 2-dim. orthonormal basis such that $\psi_1 = e_1$ and

$$\psi_2 = e^{i\lambda} s e_1 + c e_2, \quad (35)$$

with $s = \sin \alpha$, $c = \cos \alpha$ for some angle α . Now, completing the basis $\{e_1, e_2, e_3, \dots, e_k\}$ in $\mathbb{C}^k$ one easily finds that the matrix elements of $\tilde{A}$ has a form of the following direct sum

$$\tilde{A} = \left(\begin{array}{c|c} e^{-i\lambda}s & c \\ \hline -c & e^{i\lambda}s \end{array} \right) \oplus e^{-i\lambda}s \mathbb{I}_{k-2} . \quad (36)$$

Hence

$$\tilde{A}\tilde{A}^\dagger = \mathbb{I}_2 \oplus s^2 \mathbb{I}_{k-2} , \quad (37)$$

which proves (34) since all eigenvalues of $\tilde{A}\tilde{A}^\dagger - \{1, 1, s^2, \dots, s^2\}$ are bounded by 1.

Now, our new positive maps can be useful in detecting entanglement only if they are not completely positive. It is easy to check that the corresponding Choi matrix

$$W_{2k} = \sum_{i,j=1}^{2k} e_{ij} \otimes \Psi_{2k}^0(e_{ij}) , \quad (38)$$

possesses 2 negative eigenvalues $\{-1, (2-k)/k\}$ (unless $k=2$). Hence, (38) defines true entanglement witness in $\mathbb{C}^{2k} \otimes \mathbb{C}^{2k}$. As usual using Dirac notation we define $e_{kl} := |e_k\rangle\langle e_l|$. Note, that the corresponding Brauer-Hall witness possesses only one negative eigenvalue “-1”. Hence these two classes are different (unless $k=2$).

IV. PROPERTIES OF NEW ENTANGLEMENT WITNESSES

In this section we study basis properties of W_{2k} .

A. W_{2k} are indecomposable

To show that W_{2k} is indecomposable one needs to define a PPT state ρ in $\mathbb{C}^{2k} \otimes \mathbb{C}^{2k}$ such that

$\text{Tr}(W_{2k}\rho_{2k}) < 0$. Consider the following operator

$$\rho_{2k} = \sum_{i,j=1}^{2k} e_{ij} \otimes \rho_{ij}^{(2k)} , \quad (39)$$

where the $2k \times 2k$ blocks are defined as follows: diagonal blocks

$$\rho_{ii}^{(2k)} = N_k \left(\begin{array}{c|c} k\mathbb{I}_k & \mathbb{O}_k \\ \hline \mathbb{O}_k & \mathbb{I}_k \end{array} \right) , \quad (40)$$

for $i = 1, \dots, k$, and

$$\rho_{ii}^{(2k)} = N_k \left(\begin{array}{c|c} \mathbb{I}_k & \mathbb{O}_k \\ \hline \mathbb{O}_k & k\mathbb{I}_k \end{array} \right) , \quad (41)$$

for $i = k+1, \dots, 2k$. The off-diagonal blocks are form:

$$\rho_{i,i+k}^{(2k)} = -N_k W_{i,i+k}^{(2k)} , \quad (42)$$

for $i = 1, \dots, k$,

$$\rho_{ij}^{(2k)} = N_k e_{ij} , \quad (43)$$

for $i = 1, \dots, k$, $j = k+1, \dots, 2k$ and $j \neq i+k$ and

$$\rho_{ij}^{(2k)} = \mathbb{O}_k , \quad (44)$$

otherwise. The normalization factor N_k is given by

$$1/N_k = 2k^2(k+1) .$$

Direct calculation shows that

$$\rho \geq 0 , \quad \rho^\Gamma \geq 0 , \quad \text{Tr}\rho = 1 .$$

For example for $k=2$ one obtains the following density operator

$$\rho_4 = \frac{1}{24} \begin{pmatrix} 2 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 \\ \cdot & 2 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \hline \cdot & \cdot & \cdot & \cdot & 2 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & 2 & \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \hline \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ 1 & \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & 2 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 2 & \cdot & \cdot \\ \hline \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 2 & \cdot \\ 1 & \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & 2 \end{pmatrix}. \quad (45)$$

One easily finds for the trace

which proves indecomposability of W_{2k} .

In order to prove that W_{2k} is atomic one has to define a PPT state D_{2k} such that Schmidt rank of D_{2k} and of its partial transposition D_{2k}^Γ is bounded by 2 and show that $\text{Tr}(W_{2k}D_{2k}) < 0$. It is clear that atomicity implies indecomposability but for clarity of presentation we treat these two notions independently. Let us introduce the following family of product vectors

Define now the following positive operator

One easily finds for its partial transposition

Now, it is clear from that both D_{2k} and $D_{2k}^{\mathbf{r}}$ are constructed out of rank-1 projectors and Schmidt rank of each projector is 1 or 2. Therefore

Finally, one finds for the trace

which shows that W_{2k} defines atomic entanglement witness.

To show that W_{2k} is optimal we use the following result Lewenstein et. al. [7]: if the family of product vectors $\psi \otimes \phi \in \mathbb{C}^{2k} \otimes \mathbb{C}^{2k}$ satisfying

span the total Hilbert space $\mathbb{C}^{2k} \otimes \mathbb{C}^{2k}$, then W is optimal. Let us introduce the following sets of vectors:

and

$$g_{mn} = e_m + ie_n ,$$

for each $1 \leq m < n \leq 2k$. It is easy to check that $(2k)^2$ vectors $\psi_\alpha \otimes \psi_\alpha^*$ with ψ_α belonging to the set

$$\{e_l, f_{mn}, g_{mn}\} ,$$

are linearly independent and hence they do span $\mathbb{C}^{2k} \otimes \mathbb{C}^{2k}$. Direct calculation shows that

$$\langle \psi_\alpha \otimes \psi_\alpha^* | W | \psi_\alpha \otimes \psi_\alpha^* \rangle = 0 , \quad (51)$$

which proves that W_{2k} is an optimal EW.

D. W_{2k} have circulant structure

Finally, let us note that W_{2k} displays so called circulant structure [47, 48]. Let $\{e_1, \dots, e_d\}$ be an orthonormal basis in $\mathbb{C}^d$ and let $S : \mathbb{C}^d \rightarrow \mathbb{C}^d$ be a shift operator (elementary permutation) defined by

$$S e_j = e_{j+1} , \quad j = 1, \dots, d \pmod{d} . \quad (52)$$

Now, introducing

$$\Sigma_0 = \text{span}\{e_1 \otimes e_1, \dots, e_d \otimes e_d\} , \quad (53)$$

define

$$\Sigma_\alpha = (\mathbb{1} \otimes S^\alpha) \Sigma_0 , \quad \alpha = 0, \dots, d-1 . \quad (54)$$

A bipartite operator $X : \mathbb{C}^d \otimes \mathbb{C}^d \rightarrow \mathbb{C}^d \otimes \mathbb{C}^d$ displays circulant structure if

$$X = X_0 \oplus \dots \oplus X_{d-1} , \quad (55)$$

such that each X_α is supported on Σ_α . It is therefore clear that

$$X_\alpha = \sum_{i,j=1}^d x_{ij}^{(\alpha)} e_{ij} \otimes S^\alpha e_{ij} S^{\alpha\dagger} , \quad (56)$$

where $[x_{ij}^{(\alpha)}]$ is $d \otimes d$ complex matrix for each $\alpha = 0, \dots, d-1$, i.e. a circulant bipartite operator X is uniquely defined by the collection of d complex matrices $[x_{ij}^{(\alpha)}]$.

V. STRUCTURAL PHYSICAL APPROXIMATION

It is well known that positive maps cannot be directly implemented in the laboratory. The idea of *structural*

physical approximation (SPA) [42, 43] is to mix a positive map Λ with some completely positive map making the mixture $\tilde{\Lambda}$ completely positive. In the recent paper [44] the authors analyze SPA to a positive map $\Lambda : \mathcal{H}_A \rightarrow \mathcal{H}_B$ obtained through minimal admixing of white noise

$$\tilde{\Lambda}(\rho) = p \frac{\mathbb{I}_B}{d_B} \text{Tr}(\rho) + (1-p)\Lambda(\rho) . \quad (57)$$

The minimal means that the positive mixing parameter $0 < p < 1$ is the smallest one for which the resulting map $\tilde{\Lambda}$ is completely positive, i.e. it defines a quantum channel. Equivalently, one may introduce SPA of an entanglement witness W :

$$\tilde{W} = \frac{p}{d_A d_B} \mathbb{I}_A \otimes \mathbb{I}_B + (1-p)W , \quad (58)$$

where p is the smallest parameter for which $\tilde{W}$ is a positive operator in $\mathcal{H}_A \otimes \mathcal{H}_B$, i.e. it defines (possibly unnormalized) state.

It was conjectured [44] that SPA to optimal positive maps correspond to entanglement breaking maps (channels). Equivalently, SPA to optimal entanglement witnesses correspond to separable (unnormalized) states. It turns out that the family of optimal maps/witnesses constructed in this paper does support this conjecture.

The corresponding SPA of W_{2k} is given by

$$\tilde{W}_{2k} = \frac{p}{(2k)^2} \mathbb{I}_{2k} \otimes \mathbb{I}_{2k} + (1-p)W_{2k} . \quad (59)$$

Using the fact that the maximal negative eigenvalue of W_{2k} equals “−1” one easily finds the following condition for the positivity of $\tilde{W}_{2k}$

$$p \geq \frac{d}{d+1} , \quad (60)$$

with $d = 2k$. Surprisingly, one obtains the same bound for p as in Eqs. (26), (33) and (65) in [44].

Now, to show that SPA of Ψ_{2k}^0 (or equivalently W_{2k}) is entanglement breaking (equivalently separable) we use the following

Lemma 1 [49] *Let $\Lambda : \mathcal{B}(\mathbb{C}^d) \rightarrow \mathcal{B}(\mathbb{C}^d)$ be a positive unital map. Then SPA of Λ is entanglement breaking if Λ detects all entangled isotropic states in $\mathbb{C}^d \otimes \mathbb{C}^d$.*

Indeed, let

$$\rho_p = \frac{p}{d^2} \mathbb{I}_d \otimes \mathbb{I}_d + (1-p)P_d^+ , \quad (61)$$

be an isotropic state which is known to be entangled iff

$$p < \frac{1}{d+1} . \quad (62)$$

Now, assume that $(\mathbb{1} \otimes \Lambda)\rho_p$ is not positive if ρ_p is entangled. Using $\Lambda(\mathbb{I}_d) = \mathbb{I}_d$ one obtains

$$(\mathbb{1} \otimes \Lambda)\rho_p = \frac{p}{d^2} \mathbb{I}_d \otimes \mathbb{I}_d + (1-p)W, \quad (63)$$

that is, $(\mathbb{1} \otimes \Lambda)\rho_p = \widetilde{W}$. Now, if $\widetilde{W}$ is positive, then ρ_p has to be separable (otherwise it would be detected by Λ). But since $\mathbb{1} \otimes \Lambda$ sends separable states into separable states one concludes that $\widetilde{W}$ is separable (or equivalently $\widetilde{\Lambda}$ is entanglement breaking).

Lemma 2 *If in addition Λ is self-dual, i.e.*

$$\text{Tr}(A \cdot \Lambda(B)) = \text{Tr}(\Lambda(A) \cdot B), \quad (64)$$

for all $A, B \in \mathcal{B}(\mathbb{C}^d)$, then it is enough to check whether all entangled isotropic states are detected by the witness W , i.e. $\text{Tr}(W\rho_p) < 0$ for all p satisfying (62).

Again, the proof is very easy. One has

$$\begin{aligned} \text{Tr}(\rho_p \cdot W) &= \text{Tr}(\rho_p \cdot (\mathbb{1} \otimes \Lambda)P_d^+) \\ &= \text{Tr}((\mathbb{1} \otimes \Lambda)\rho_p \cdot P_d^+), \end{aligned} \quad (65)$$

where in the last equality we used the self-duality of Λ . Now, if $\text{Tr}(\rho_p \cdot W) < 0$ for p satisfying (62), then $(\mathbb{1} \otimes \Lambda)\rho_p$ is not positive (otherwise its trace with the projector P_d^+ would be positive). Hence by Lemma 1 SPA $\widetilde{\Lambda}$ is entanglement breaking.

We are prepared to show that SPA for Ψ_{2k}^0 is entanglement breaking.

Lemma 3 Ψ_{2k}^0 *is self-dual.*

One checks by direct calculations that

$$\text{Tr}(e_{kl} \cdot \Psi_{2k}^0(e_{mn})) = \text{Tr}(\Psi_{2k}^0(e_{kl}) \cdot e_{mn}), \quad (66)$$

for all $k, l, m, n = 1, \dots, 2k$. Hence, due to the Lemma 2, to show that SPA for Ψ_{2k}^0 is entanglement breaking is it enough to prove

Lemma 4 $\text{Tr}(W_{2k}\rho_p) < 0$ *for all p satisfying (62).*

To prove it let us note that

$$\text{Tr}(W_{2k}\rho_p) = \frac{p}{(2k)^2} \text{Tr} W_{2k} + (1-p) \text{Tr}(W_{2k}P_{2k}^+).$$

Now, $\text{Tr} W_{2k} = 2k$, and

$$\text{Tr}(W_{2k}P_{2k}^+) = \frac{1}{2k} \sum_{m,n=1}^{2k} \langle m | \Psi_{2k}^0(e_{mn}) | n \rangle.$$

Finally, using definition of Ψ_{2k}^0 one gets

$$\sum_{m,n=1}^{2k} \langle m | \Psi_{2k}^0(e_{mn}) | n \rangle = -2k,$$

and hence

$$\text{Tr}(W_{2k}\rho_p) = \frac{p(d+1)-1}{d}, \quad (67)$$

with $d = 2k$. Now, if ρ_p is entangled, i.e. $p < 1/(d+1)$, then $\text{Tr}(W_{2k}\rho_p) < 0$ which shows that W_{2k} detects all entangled isotropic states.

VI. CONCLUSIONS

We have provided a new construction of EWs in $\mathbb{C}^d \otimes \mathbb{C}^d$ with $d = 2k$. It was shown that these EWs are indecomposable, i.e. they are able to detect PPT entangled state. Moreover, they are so called atomic EWs, i.e. they are able to detect PPT entangled states ρ such that both ρ and ρ^Γ possess Schmidt number 2. The crucial property of our witnesses is they optimality, i.e. they are no other witnesses which can detect more entangled states.

Equivalently, our construction gives rise to the new class of positive maps in algebras of $d \times d$ complex matrices. For $k = 2$ this construction reproduces old example of Robertson map [28] and hence [39] defines the special case of Brauer-Hall maps [36, 37].

Let us observe that if $\Lambda : \mathcal{B}(\mathbb{C}^d) \rightarrow \mathcal{B}(\mathbb{C}^d)$ is a positive indecomposable map then for any unitaries $U_1, U_2 : \mathbb{C}^d \rightarrow \mathbb{C}^d$ a new map

$$\Lambda^{U_1 U_2}(A) := U_1 \Lambda(U_2^\dagger A U_2) U_1^\dagger, \quad (68)$$

is again positive and indecomposable [39]. This observation enables one to generalize Robertson map Φ_{2k}^0 and our new map Ψ_{2k}^0 to $\Phi_{2k}^{U_1 U_2}$ and $\Psi_{2k}^{U_1 U_2}$. Note, that if $U_1 = U_2 = U$ given by (15), then $\Phi_{2k}^U := \Phi_{2k}^{UU}$ defines Breuer-Hall map. Therefore, $\Psi_{2k}^U := \Psi_{2k}^{UU}$ may be regarded as a Breuer-Hall-like generalization of our primary map Ψ_{2k}^0 .

It should be stressed, an EW defined by the Breuer-Hall map and EW W_{2k} introduced in this paper are different, i.e. they do detect different classes of PPT entangled states. Direct calculation shows that the PPT entangled state (39) is not detected by the Breuer-Hall witness. On the other hand consider the family of PPT entangled state introduced in [36]

$$\rho(\lambda) = \lambda P_d^+ + (1-\lambda)\rho_0, \quad (69)$$

with

$$\rho_0 = \frac{2}{d(d+1)} U_0 P_S U_0^\dagger, \quad (70)$$

and P_S being the projector onto the subspace of states symmetric under the swap operation. It was shown that $\rho(\lambda)$ is PPT for $0 \leq \lambda \leq 1/(d+2)$. Moreover,

Breuer-Hall witness detects all entangled states within λ -family (69) (both PPT and NPT). Direct calculation shows that our witness W_{2k} does not detect PPT entangled states in (69).

Interestingly, the partial transposition W_{2k}^Γ defines an EW with $k(k-1)/2$ negative eigenvalues (all equal to -1). For $k = 2$ it gives exactly one negative eigenvalue (the fact well known from the family of Breuer-Hall maps in 4 dimensions). Therefore, this example provide an EW with multiple negative eigenvalues. However, contrary to the Breuer-Hall maps we were not able to show that W_{2k}^Γ is optimal.

We have shown that structural physical approximation for our new class of positive maps gives rise to entanglement breaking channels. This result supports recent conjecture by Korbicz et. al. [44].

Finally, let us mention some open questions. In

this paper we have used reduction map as a building block to construct other optimal maps. Can we use other positive maps as building blocks? Is it true that properties of building blocks (like optimality and/or atomicity) are shared by the map which is built out of them?

Acknowledgement

We thank Antonio Acin and Andrzej Kossakowski for valuable discussions. This work was partially supported by the Polish Ministry of Science and Higher Education Grant No 3004/B/H03/2007/33 and by the Polish Research Network *Laboratory of Physical Foundations of Information Processing*.

-
- [1] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information*, (Cambridge University Press, Cambridge, England, 2000).
 - [2] R. Horodecki, P. Horodecki, M. Horodecki and K. Horodecki, *Rev. Mod. Phys.* **81**, 865 (2009).
 - [3] M. Horodecki, P. Horodecki and R. Horodecki, *Phys. Lett. A* **223**, 1 (1996).
 - [4] B.M. Terhal, *Phys. Lett. A* **271**, 319 (2000).
 - [5] B. M. Terhal, *Theor. Comput. Sci.* **287**, 313 (2002).
 - [6] M. Lewenstein, B. Kraus, J. I. Cirac, and P. Horodecki, *Phys. Rev. A* **62**, 052310 (2000).
 - [7] M. Lewenstein, B. Kraus, P. Horodecki, and J. I. Cirac, *Phys. Rev. A* **63**, 044304 (2001).
 - [8] B. Kraus, M. Lewenstein, and J. I. Cirac, *Phys. Rev. A* **65**, 042327 (2002).
 - [9] P. Hyllus, O. Gühne, D. Bruss, and M. Lewenstein *Phys. Rev. A* **72**, 012321 (2005).
 - [10] D. Bruß, *J. Math. Phys.* **43**, 4237 (2002).
 - [11] G. Tóth and O. Gühne, *Phys. Rev. Lett.* **94**, 060501 (2005)
 - [12] R. A. Bertlmann, H. Narnhofer and W. Thirring, *Phys. Rev. A* **66**, 032319 (2002).
 - [13] F.G.S.L. Brandão, *Phys. Rev. A* **72**, 022310 (2005).
 - [14] G. Sarbicki, *J. Phys. A: Math. Theor.* **41**, 375303 (2008).
 - [15] D. Chruściński and A. Kossakowski, *J. Phys. A: Math. Theor.* **41**, 145301 (2008).
 - [16] M. Bourennane, M. Eibl, C. Kurtsiefer, S. Gaertner, H. Weinfurter, O. Gühne, P. Hyllus, D. Bruss, M. Lewenstein and A. Sanpera, *Phys. Rev. Lett.* **92**, 087902 (2004).
 - [17] L.-A. Wu, S. Bandyopadhyay, M. S. Sarandy, and D. A. Lidar, *Phys. Rev. A* **72**, 032309 (2005).
 - [18] A.C. Doherty, P.A. Parrilo and F.M. Spedalieri, *Phys. Rev. Lett.* **88**, 187904 (2002).
 - [19] F.G.S.L. Brandão and R.O. Vianna, *Phys. Rev. Lett.* **93**, 220503 (2004).
 - [20] J. Eisert, P. Hyllus, O. Gühne, and M. Curty, *Phys. Rev. A* **70**, 062317 (2004);
 - [21] P. Hyllus, O. Guehne, D. Bruß, M. Lewenstein, *Phys. Rev. A* **72**, 012321 (2005).
 - [22] M.-D. Choi, *Lin. Alg. Appl.* **10**, 285 (1975).
 - [23] A. Jamiolkowski, *Rep. Math. Phys.* **3**, 275 (1972).
 - [24] S.L. Woronowicz, *Rep. Math. Phys.* **10**, 165 (1976).
 - [25] E. Størmer, *Acta Math.* **110**, 233 (1963).
 - [26] M.-D. Choi, *Lin. Alg. Appl.* **12**, 95 (1975); M.-D. Choi, *J. Operator Theory*, **4**, 271 (1980).
 - [27] S.L. Woronowicz, *Comm. Math. Phys.* **51**, 243 (1976).
 - [28] A.G. Robertson, *Math. Proc. Camb. Phil. Soc.*, **94**, 71 (1983). **34**, 87 (1983); A.G. Robertson, *J. London Math. Soc. (2)* **32**, 133 (1985).
 - [29] W.-S. Tang, *Lin. Alg. Appl.* **79**, 33 (1986).
 - [30] K. Tanahashi and J. Tomiyama, *Canad. Math. Bull.* **31**, 308 (1988).
 - [31] H. Osaka, *Lin. Alg. Appl.* **153**, 73 (1991); *ibid.* **186**, 45 (1993).
 - [32] F. Benatti, R. Floreanini and M. Piani, *Phys. Lett. A* **326**, 187 (2004).
 - [33] K.-C. Ha, *Publ. RIMS, Kyoto Univ.* **34**, 591 (1998)
 - [34] K.-C. Ha and S.-H. Kye, *J. Phys. A: Math. Gen.* **38**, 9039 (2005); *Phys. Lett. A* **325**, 315 (2004).
 - [35] A. Kossakowski, *Open Sys. Information Dyn.* **10**, 213 (2003).
 - [36] H.-P. Breuer, *Phys. Rev. Lett.* **97**, 0805001 (2006).
 - [37] W. Hall, *J. Phys. A: Math. Gen.* **39**, (2006) 14119.
 - [38] D. Chruściński and A. Kossakowski, *Open Systems and Inf. Dynamics*, **14**, 275 (2007).
 - [39] D. Chruściński and A. Kossakowski, *J. Phys. A: Math. Theor.* **41**, 215201 (2008).
 - [40] D. Chruściński and A. Kossakowski, *Spectral conditions for positive maps*, arXiv:0809.4909 (to be published in *Comm. Math. Phys.*)
 - [41] D. Chruściński and A. Kossakowski, *Phys. Lett. A* **373**, 2301 (2009).
 - [42] P. Horodecki, *Phys. Rev. A* **68**, 052101 (2003).
 - [43] P. Horodecki and A. Ekkert, *Phys. Rev. Lett.* **89**,

- 127902 (2002).
- [44] J.K. Korbicz, M.L. Almeida, J. Bae, M. Lewenstein and A. Acin, Phys. Rev. A **78**, 062105 (2008).
 - [45] B. Terhal and P. Horodecki, Phys. Rev. A **61**, 040301 (2000); A. Sanpera, D. Bruss and M. Lewenstein, Phys. Rev. A **63**, 050301(R) (2001)
 - [46] M. Horodecki and P. Horodecki, Phys. Rev. **A** 59, 4206 (1999).
 - [47] D. Chruściński and A Kossakowski, Phys. Rev. A **74**, 022308 (2006)
 - [48] D. Chruściński and A Kossakowski, Phys. Rev. A **76**, 032308 (2007); D. Chruściński and A. Pittenger, J. Phys. A: Math. Theor. **41**, 385301 (2008).
 - [49] We thank Antonio Acin for his remarks.